Math-2 Lesson 9-3

Exponential Function


The "Parent" Exponential Function $y = b_{base}^{x}$


- $y = 2^{x}$ (base 2 exponential function)
- $y = 3^{\chi}$ (base 3 exponential function)
- $y = \left(\frac{1}{2}\right)^x$ (base 1/2 exponential function)

The base MUST BE positive and CANNOT equal 1.

$$b = (0,1) \cup (1,\infty)$$


Fill in the output values of the table and graph the points.








Exponential Function $f(x) = 2^x$


Will the '<u>y' value ever reach zero</u> (on the left end of the graph)? As the denominator gets bigger and bigger, the decimal version of the fraction gets smaller and smaller.















Negative Exponent Property




$$f(x) = 2^x$$
 k(x) = $2^x + 4$

X	2()	f(x)	k(x)
-2	2^{-2}	0.25	4.25
-1	2^{-1}	0.5	4.5
0	2^{0}	1	5
1	2^{1}	2	6
2	2^2	4	8


Horizontal 3 asymptote: 3

$$y = 0$$
$$y = 4$$

Domain =?
$$\begin{array}{l} x = (-\infty, \infty) \\ x = (-\infty, \infty) \end{array}$$



Transformations of the Exponential Function



Summary

k = 0

 $g(x) = ab^x + k$ 1) Start with

2) Find the value of 'k' (horizontal asymptote).

$$g(x) = ab^x + k \quad \rightarrow$$


3) Substitute the y-intercept


$$(0, 1) \rightarrow y = ab^{x} \rightarrow 1 = ab$$

$$\rightarrow a = 1 \rightarrow y = b^{x}$$

4) Substitute a "nice" x-y pair from the graph into the equation.

$$(1, 2) \rightarrow y = b^x \rightarrow 2 = b^1 \rightarrow b = 2 \rightarrow y = 2^x$$





$$(1, 5) \rightarrow 5 = b^1 + 3 \rightarrow b = 2$$
$$y = 2^x + 3$$

