

Math-3A

Lesson 11-10

3rd Degree Polynomials

Find the zeroes of the following 3rd degree Polynomial

$$y = x^3 + 5x^2 + 4x \quad \text{Set } y = 0$$

$$0 = x^3 + 5x^2 + 4x \quad \text{Factor out the common factor.}$$

$$0 = x(x^2 + 5x + 4) \quad \text{Factor the quadratic}$$

$$0 = x(x + 1)(x + 4) \quad \text{Identify the zeroes}$$

0, -1, -4

“Nice” (factorable) 3rd Degree Polynomials

$$y = ax^3 + bx^2 + cx + d$$

If it has no constant term, it will look like this:

$$y = ax^3 + bx^2 + cx$$

This can easily be factored (by taking out the common factor ‘x’).

$$y = x(ax^2 + bx + c)$$

Resulting in ‘x’ times a quadratic factor.

We have been factoring quadratics for quite a while!

“Nice” 3rd Degree Polynomial (with no constant term)

$$y = x^3 + 6x^2 + 4x + 0$$

$$0 = x^3 + 6x^2 + 4x$$

It has no constant term so it can easily be factored

into ‘x’ times a quadratic factor. $0 = x(x^2 + 6x + 4)$

$$x = 0$$

What if the quadratic factor is not factorable?

$$x = \frac{-b}{2a} = \frac{-(6)}{2(1)}$$

$$x = -3$$

$$y = f(-3) = (-3)^2 + 6(-3) + 4$$

$$y = -5$$

Convert the quadratic factor into vertex form and solve.

$$0 = (x + 3)^2 - 5$$

$$x = -3 \pm \sqrt{5}$$

Zeroes:

$$x = 0, -3 \pm \sqrt{5}$$

Factor the following “nice” 3rd degree polynomials then find the “zeroes” of the polynomial.

$$y = x^3 + 5x^2 - 14x$$

$$0 = x^3 + 5x^2 - 14x$$

$$0 = x(x^2 + 5x - 14)$$

$$0 = x(x + 7)(x - 2)$$

0, -7, 2

$$y = 3x^3 - 24x^2 + 6x$$

$$0 = 3x(x^2 - 8x + 2)$$

$$x = 0$$

$$x = \frac{-b}{2a} = \frac{-(-6)}{2(1)}$$

$$x = 4$$

$$y = f(4) = (4)^2 - 8(4) + 2$$


$$y = -14 \quad 0 = (x + 4)^2 - 14$$

$$x = -4 \pm \sqrt{14}$$

Another “Nice” 3rd Degree Polynomial

$$y = ax^3 + bx^2 + cx + d$$

This has the constant term, but it has a very useful feature:


$$y = 1x^3 + 2x^2 + 2x + 4$$

What pattern do you see?

$$3^{rd} / 1^{st} = \frac{2}{1} \quad 4^{th} / 2^{nd} = \frac{4}{2} = \frac{2}{1}$$

An easy method is “box factoring” if it has this nice pattern.

$$y = 1x^3 + 2x^2 + 2x + 4$$

These 4 terms are the numbers in the box.

Find the common factor of the 1st row.

Fill in the rest of the box.

	x	2
x^2	x^3	$2x^2$
2	$2x$	4

Rewrite in intercept form.

$$y = 1x^3 + 2x^2 + 2x + 4$$

$$y = (x^2 + 2)(x + 2)$$

Find the “zeroes.”

$$0 = (x^2 + 2)(x + 2)$$

$$0 = x^2 + 2$$

$$0 = x + 2$$

$$-2 = x^2$$

$$x = -2$$

$$x = \pm i\sqrt{2}$$

Find the zeroes using “box factoring”

$$y = 4x^3 - 5x^2 + 12x - 15$$

$$0 = (x^2 + 3)(4x - 5)$$

$$0 = x^2 + 3 \quad 4x - 5 = 0$$

$$-3 = x^2$$

$$x = \pm i\sqrt{3}$$

$$4x = 5$$

$$x = 5/4$$

	$4x$	-5
x^2	$4x^3$	$-5x^2$
3	$12x$	-15


$$x = i\sqrt{3}, -i\sqrt{3}, 5/4$$

What have we learned so far?

“Nice” Common Factor 3rd degree polynomial:

$$\begin{aligned}y &= x^3 + 3x^2 + 2x &= x(x^2 + 3x + 2) \\ & &= x(x + 1)(x + 2)\end{aligned}$$

“Nice” Factor by box 3rd degree polynomial:

$$y = 2x^3 + 3x^2 + 4x + 6$$


“Nice” Difference of Squares (of higher degree):

$$y = x^4 - 81 \quad \text{Use “m” substitution} \quad \text{Let } m^2 = x^4$$

$$y = m^2 - 81 \quad \text{Then } m = x^2$$

$$y = (m + 9)(m - 9)$$

Use “m” substitution

$$y = (x^2 + 9)(x^2 - 9)$$

$$y = (x + 3)(x - 3)(x + 3i)(x - 3i)$$

Find the zeroes. $x = -3, 3, -3i, 3i$

Convert to standard form:

$$y = (x - 3)(x^2 + 3x + 9)$$

$$y = x^3 - 27$$

There are NO x^2 terms
and NO 'x' terms

The Difference of cubes: factors
as the cubed root of each term
multiplied by a 2nd degree
polynomial. $y = x^3 - 8$

$$y = (x - 2)(ax^2 + bx + c)$$

$$y = (x - 2)(x^2 + 2x + 4)$$

	x	-3
x^2	x^3	$-3x^2$
$3x$	$3x^2$	$-9x$
9	$9x$	-27

$0x^2$ $0x$

	x	-2
x^2	x^3	$-2x^2$
$2x$	$2x^2$	$-4x$
4	$4x$	-8

$0x^2$ $0x$

The Sum of cubes: factors as the cubed root of each term multiplied by a 2nd degree polynomial. $y = x^3 + 64$

$$y = (x + 4)(ax^2 + bx + c)$$

	x	4
x^2	x^3	$4x^2$
$-4x$	$-4x^2$	$-16x$
16	$16x$	64

$0x^2$ $0x$

$$y = (x + 4)(x^2 - 4x + 16)$$