

# Math-3A

## Lesson 3-4

### Factoring “NICE” 3rd Degree Polynomials

Find the zeroes of the following 3<sup>rd</sup> degree Polynomial

$$y = x^3 + 5x^2 + 4x \quad \text{Set } y = 0$$

$$0 = x^3 + 5x^2 + 4x \quad \text{Factor out the common factor.}$$

$$0 = x(x^2 + 5x + 4) \quad \text{Factor the quadratic}$$

$$0 = x(x + 1)(x + 4) \quad \text{Identify the zeroes}$$

**0, -1, -4**

## “Nice” (factorable) 3<sup>rd</sup> Degree Polynomials

$$y = ax^3 + bx^2 + cx + \textcircled{d}$$

If it has no constant term, it will look like this:

$$y = ax^3 + bx^2 + cx$$

This can easily be factored (by taking out the common factor ‘x’).

$$y = x \textcircled{(ax^2 + bx + c)}$$

Resulting in ‘x’ times a quadratic factor.

We have been factoring quadratics for quite a while!

## “Nice” 3<sup>rd</sup> Degree Polynomial (with no constant term)

$$y = x^3 + 5x^2 + 4x \quad \bigcirc$$

It has no constant term so it can easily be factored into ‘x’ times a quadratic factor.

$$y = x(x^2 + 5x + 4)$$

***if*** the quadratic factor is “nice” we can factor that into 2 binomials.

$$y = x(x + 1)(x + 4)$$

This is now “intercept form” so we can “read off” the x-intercepts. What are they?

**0, -1, -4**

## “Nice” 3<sup>rd</sup> Degree Polynomial (with no constant term)

$$y = x^3 + 6x^2 + 4x + 0$$

$$0 = x^3 + 6x^2 + 4x$$

It has no constant term so it can easily be factored

into ‘x’ times a quadratic factor.  $0 = x(x^2 + 6x + 4)$

$$x = 0$$

**What if** the quadratic factor is not factorable?

$$x = \frac{-b}{2a} = \frac{-(6)}{2(1)}$$

$$x = -3$$

$$y = f(-3) = (-3)^2 + 6(-3) + 4$$

$$y = -5$$

Convert the quadratic factor into vertex form and solve.

$$0 = (x + 3)^2 - 5$$

$$x = -3 \pm \sqrt{5}$$

Zeroes:

$$x = 0, -3 \pm \sqrt{5}$$

Factor the following “nice” 3<sup>rd</sup> degree polynomials then find the “zeroes” of the polynomial.

$$y = x^3 + 5x^2 - 14x$$

$$0 = x^3 + 5x^2 - 14x$$

$$0 = x(x^2 + 5x - 14)$$

$$0 = x(x + 7)(x - 2)$$

0, -7, 2

$$y = 3x^3 - 24x^2 + 6$$

$$0 = 3x(x^2 - 8x + 2)$$

$$x = 0$$

$$x = \frac{-b}{2a} = \frac{-(-6)}{2(1)}$$

$$x = 4$$

$$y = f(4) = (4)^2 - 8(4) + 2$$


$$y = -14 \quad 0 = (x + 4)^2 - 14$$

$$x = -4 \pm \sqrt{14}$$

Another “Nice” 3<sup>rd</sup> Degree Polynomial

$$y = ax^3 + bx^2 + cx + d$$


This has the constant term, but it has a very useful feature:


$$y = 1x^3 + 2x^2 + 2x + 4$$

What pattern do you see?

$$3^{rd} / 1^{st} = \frac{2}{1} \quad 4^{th} / 2^{nd} = \frac{4}{2} = \frac{2}{1}$$

“factor by grouping” if it has this nice pattern.

$$y = 1x^3 + 2x^2 + 2x + 4$$
The equation  $y = 1x^3 + 2x^2 + 2x + 4$  is shown. Two red curved arrows are drawn above the equation. One arrow starts above the first term  $1x^3$  and points to the second term  $2x^2$ . The other arrow starts below the third term  $2x$  and points to the fourth term  $4$ . This indicates that the first and second terms are grouped together, and the third and fourth terms are grouped together.


Group the 1<sup>st</sup> and last pair of terms with parentheses.

$$y = (x^3 + 2x^2) + (2x + 4)$$

Factor out the common term from the first group.

$$y = x^2(x + 2) + (2x + 4)$$

Factor out the common term from the last group.

$$y = x^2(x + 2) + 2(x + 2)$$
The equation  $y = x^2(x + 2) + 2(x + 2)$  is shown. Two red ovals are drawn around the terms  $(x + 2)$  in the first and second terms, highlighting the common factor.

Common factor of  $(x + 2)$



$$y = x^2(x + 2) + 2(x + 2)$$

These two binomials are now common factors of

$$x^2 \text{ and } 2$$

Factor out the common binomial term.

$$y = (x + 2)(x^2 + 2)$$

Apply the Complex Conjugates Theorem

$$x = -2, i\sqrt{2}, -i\sqrt{2}$$

An easier method is “box factoring” if it has this nice pattern.

$$y = 1x^3 + 2x^2 + 2x + 4$$

These 4 terms are the numbers in the box.

Find the common factor of the 1<sup>st</sup> row.

Fill in the rest of the box.

	$x$	$2$
$x^2$	$x^3$	$2x^2$
$2$	$2x$	$4$

Rewrite in intercept form.

$$y = 1x^3 + 2x^2 + 2x + 4$$

$$y = (x^2 + 2)(x + 2)$$

Find the “zeroes.”

$$0 = (x^2 + 2)(x + 2)$$

$$0 = x^2 + 2$$


$$0 = x + 2$$

$$-2 = x^2$$

$$x = -2$$

$$x = \pm i\sqrt{2}$$

Which of the following 3<sup>rd</sup> degree polynomials have the “nice pattern” we saw on the previous slide?

$$y = 2x^3 + 3x^2 + 4x + 6 \quad 3^{\text{rd}} / 1^{\text{st}} = 4^{\text{th}} / 2^{\text{nd}} = 2$$


$$y = 4x^3 - 5x^2 + 12x - 15 \quad 3^{\text{rd}} / 1^{\text{st}} = 4^{\text{th}} / 2^{\text{nd}} = 3$$

$$y = 4x^3 + 8x^2 + 5x + 10 \quad 3^{\text{rd}} / 1^{\text{st}} = 4^{\text{th}} / 2^{\text{nd}} = 5/4$$

$$y = -6x^3 + 7x^2 - 18x + 21 \quad 3^{\text{rd}} / 1^{\text{st}} = 4^{\text{th}} / 2^{\text{nd}} = 3$$

All of them!

Find the zeroes using “box factoring”

$$y = 4x^3 - 5x^2 + 12x - 15$$

$$0 = (x^2 + 3)(4x - 5)$$

$$0 = x^2 + 3$$

$$4x - 5 = 0$$

$$-3 = x^2$$

$$4x = 5$$

$$x = \pm i\sqrt{3}$$

$$x = 5/4$$

	$4x$	$-5$
$x^2$	$4x^3$	$-5x^2$
$3$	$12x$	$-15$


$$x = i\sqrt{3}, -i\sqrt{3}, 5/4$$

What have we learned so far?

“Nice” Common Factor 3<sup>rd</sup> degree polynomial:

$$\begin{aligned}y &= x^3 + 3x^2 + 2x &= x(x^2 + 3x + 2) \\ & &= x(x + 1)(x + 2)\end{aligned}$$

“Nice” Factor by box 3<sup>rd</sup> degree polynomial:

$$y = 2x^3 + 3x^2 + 4x + 6$$


“Nice” Difference of Squares (of higher degree):

$$y = x^4 - 81 \quad \text{Use “m” substitution} \quad \text{Let } m^2 = x^4$$

$$y = m^2 - 81$$

$$\text{Then } m = x^2$$

$$y = (m + 9)(m - 9)$$

Use “m” substitution

$$y = (x^2 + 9)(x^2 - 9)$$

$$y = (x + 3)(x - 3)(x + 3i)(x - 3i)$$

Find the zeroes.  $x = -3, 3, -3i, 3i$

Convert to standard form:

$$y = (x - 3)(x^2 + 3x + 9)$$

$$y = x^3 - 27$$

There are NO  $x^2$  terms  
and NO 'x' terms

The Difference of cubes: factors  
as the cubed root of each term  
multiplied by a 2<sup>nd</sup> degree  
polynomial.  $y = x^3 - 8$

$$y = (x - 2)(ax^2 + bx + c)$$

$$y = (x - 2)(x^2 + 2x + 4)$$

	$x$	$-3$
$x^2$	$x^3$	$-3x^2$
$3x$	$3x^2$	$-9x$
$9$	$9x$	$-27$

$0x^2$     $0x$

	$x$	$-2$
$x^2$	$x^3$	$-2x^2$
$2x$	$2x^2$	$-4x$
$4$	$4x$	$-8$

$0x^2$     $0x$

The Sum of cubes: factors as the cubed root of each term multiplied by a 2<sup>nd</sup> degree polynomial.  $y = x^3 + 64$

$$y = (x + 4)(ax^2 + bx + c)$$

	$x$	$4$
$x^2$	$x^3$	$4x^2$
$-4x$	$-4x^2$	$-16x$
$16$	$16x$	$64$

$0x^2$     $0x$

$$y = (x + 4)(x^2 - 4x + 16)$$